

## JOURNAL PUBLICATIONS

A Prodigiosin Analogue Inactivates NADPH Oxidase in Macrophage Cells by Inhibiting Assembly of P47phox and Rac.

Nakashima T., Iwashita T., Fujita T., Sato E., Niwano Y., Kohno M., Kuwahara S., Harada N., Takeshita S., and Oda T.

*J. Biochem.* **143**, 107-115 (2008)

Overexpression of Post-Translationally Modified Peptides in *Escherichia coli* by Co-Expression with Modifying Enzymes.

Sugase K., Landes M. A., Wright P. E., and Martinez-Yamout M.

*Protein Expres. Purif.* **57**, 108-115 (2008)\*

Cytotoxic Action Mode of a Novel Porphyrin Derivative Isolated from Harmful Red Tide Dinoflagellate *Heterocapsa circularisquama*.

Kim D., Miyazaki Y., Nakashima T., Iwashita T., Fujita T., Yamaguchi K., Choi K. S., and Oda T.

*J. Biochem. Mol. Toxicol.* **22**, 158-165 (2008)

Sequential Glucosylation of a Furofuran Lignan, (+)-Sesaminol, by *Sesamum indicum* UGT71A9 and UGT94D1 Glucosyltransferases.

Noguchi A., Fukui Y., Iuchi-Okada A., Kakutani S., Satake H., Iwashita T., Nakao M., Umezawa T., Ono E.

*Plant J.* **54**, 415-427 (2008)\*

Characterization of Cd Translocation and Identification of the Cd Form in Xylem Sap of the Cd-Hyperaccumulator *Arabidopsis halleri*.

Ueno D., Iwashita T., Zhao F. J., and Ma J. F.

*Plant Cell Physiol.* **49**, 540-548 (2008)

Interaction of Lipopolysaccharide and Phospholipid in Mixed Membranes: Solid-State <sup>31</sup>P-NMR Spectroscopic and Microscopic Investigations.

Nomura K., Inaba T., Morigaki K., Brandenburg K., Seydel U., and Kusumoto S.

*Biophys. J.* **95**, 1226-1238 (2008)\*

Orientation of Fluorinated Cholesterol in Lipid Bilayers Analyzed by <sup>19</sup>F Tensor Calculation and Solid-State NMR.

Matsumori N., Kasai Y., Oishi T., Murata M., and Nomura K.

*J. Am. Chem. Soc.* **130**, 4757-4766 (2008)

Inconspicamide, New N-Acylated Serinol from the Marine Sponge *Stelletta inconspicua*.  
Ueoka R., Fujita T., Iwashita T., van Soest R. W., and Matsunaga S.  
*Biosci. Biotechnol. Biochem.* **72**, 3055-3058 (2008)

Rapid and Efficient Synthesis of Dysiherbaine and Analogues to Explore Structure-Activity Relationships.  
Sasaki M., Tsubone K., Aoki K., Akiyama N., Shoji M., Oikawa M., Sakai R., and Shimamoto K.  
*J. Org. Chem.* **73**, 264-273 (2008)

A Positive Charge at the N-Terminal Segment of Discrepin Increases the Blocking Effect of K<sup>+</sup> Channels Responsible for the IA Currents in Cerebellum Granular Cells.  
Rameo S., Corzo G., Vasile A., Satake H., Prestipino G., and Possani L. D.  
*Biochem. Biophys. Acta.* **780**, 750-755 (2008)

TMG-Chitotriomycin, an Enzyme Inhibitor Specific for Insect and Fungal β-N-Acetyl-glucosaminidases, Produced by Actinomycete *Streptomyces anulatus* NBRC 13369.  
Usuki H., Nitoda T., Ichikawa M., Yamaji N., Iwashita T., Komura H., and Kanzaki H.  
*J. Amer. Chem. Soc.* **130**, 4146-4152 (2008)

Low-Affinity Excitatory Amino Acid Uptake in Hippocampal Astrocytes: A Possible Role of Na<sup>+</sup>/Dicarboxylate Cotransporters.  
Holten A. T., Danbolt N. C., Shimamoto K., and Gundersen V.  
*Glia* **56**, 990-997 (2008)

Crystal Structure of Squid Rhodopsin with Intracellularly Extended Cytoplasmic Region.  
Shimamura T., Hiraki K., Takahashi N., Hori T., Ago H., Masuda K., Takio K., Ishiguro M., and Miyano M.  
*J. Biol. Chem.* **283**, 17753-17756 (2008)\*

A Novel Biological Role of Tachykinins as an Upregulator of Oocyte Growth: Identification of an Evolutionary Origin of Tachykinergic Functions in the Ovary of the Ascidian *Ciona intestinalis*.  
Aoyama M., Kawada T., Fujie M, Hotta K. Sakai T., Sekiguchi T., Oka K., Satoh N., and Satake H.  
*Endocrinology* **149**, 4346-4356 (2008)\*

Microstructure of Poly(γ-Glutamic acid) Produced by *Bacillus subtilis* Consisting of Clusters of D- and L-Glutamic Acid Repeating Units.  
Wang F., Ishiguro M., Mutsukado M., Fujita K., and Tanaka T.  
*J. Agric. Food Chem.* **56**, 4225-4228 (2008)\*

Characterization of a Novel Vasopressin/Oxytocin Superfamily Peptide and its Receptor from an Ascidian, *Ciona intestinalis*.

Kawada T., Sekiguchi T., Itoh Y., Ogasawara M., and Satake H.  
*Peptides* **29**, 1672-1678 (2008)\*

A Common "Hot Spot" Confers hERG Blockade Activity to Alpha-Scorpion Toxins Affecting K<sup>+</sup> Channels.

Abdel-Mottaleb Y., Corzo G., Martin-Eauclaire M.F., Satake H., Céard B., Peigneur S., Nambaru P., Bougis P.E., Possani L. D., and Tytgat J.  
*Biochem. Pharmacol.* **76**, 805-815 (2008)

Neuropeptide Receptor Transcriptome Reveals Unidentified Neuroendocrine Pathways.

Yamanaka N., Yamamoto S., Zitnan D., Watanabe K., Kawada T., Satake H., Kaneko Y., Hinuma K., Tanaka Y., Shinoda T., and Kataoka H.  
*PLoS ONE* **3**, e3048 (2008)

Novel Ubiquitous Promoters and Expression-Vector Optimization in Ascidian Embryos.

Shimai K., Hirano A., Kitaura Y., Kitano Y., Itoh A., Kiuchi A., Sasaki N., and Nishikata T.  
*Invertebr. Reprod. Dev.* **51**, 103-110 (2008)

## BOOKS AND REVIEWS

Recent Advances and Perceptions in Studies of Heterodimerization Between G Protein-Coupled Receptors.

Satake H., and Sakai T.

*Protein Pept. Lett.* **15**, 300-308 (2008)\*

Neuropeptides and their Functions in *Hydra*.

Takahashi T., Hayakawa E., Koizumi O., and Fujisawa T.

*Acta Biol. Hung.* **59** (suppl), 227-235 (2008)

Glutamate Transporter Blockers for Elucidation of the Function of Excitatory Neurotransmission Systems.

Shimamoto K.

*Chem. Rec.*, **8**, 182-199 (2008)\*

Improved Application of Semiconducting Metal Oxides as a Detector for High-Resolution Gas Chromatography

Komura H.

ACS Symposium Series 988, "Food Flavor: Chemistry, Sensory Evaluation, and Biological Activity", pp. 36-44, Eds.

Tamura H., Ebeler S.E., Kubota K., and Takeoka G.R. American Chemical Society, Washington DC (2008)\*

A Specific Transporter for Iron (III)–Phytosiderophore Complex Involved in Iron Acquisition by Barley Roots.  
Murata Y., Harada E., Sugase K., Namba K., Horikawa M., Ma J.F., Yamaji N., UenoD., Nomoto K., Iwashita T.,  
Kusumoto S.  
*Pure Appl. Chem.* **80**, 2689-2697 (2008)\*

Analysis of Cyclodextrins and their Derivatives by Fourier Transform Ion Cyclotron Resonance Mass Spectrometry with an Ultra-High 9.4T Magnetic Field: Mixture of Beta-Cyclodextrin and its Aminated Molecules.  
Yamagaki T.  
*Proceedings of the 14th International Cyclodextrins Symposium* 81-84 (2008)